

FOOT-AND-MOUTH DISEASE + HAEMORRHAGIC SEPTICAEMIA VACCINE



COMPOSITION

Each dose of FH VAC® combined vaccine contains aziridine inactivated antigens from BHK cell culture grown foot-and-mouth disease virus serotypes O (strain IND/R2/75), A (strain IND/40/2000) and Asia1 (strain IND/63/72), and formalin killed cultures of *Pasteurella multocida* (strain P52), adjuvanted with mineral oil. Thiomersal ($\leq 0.02\%$ w/v) is added as preservative.

DOSAGE & ADMINISTRATION

2 mL in cattle and buffaloes,
1 mL in sheep and goats.
For deep intramuscular use only.

Presentation
Pack: 30 mL | 100 mL



FH VAC®

एफएच वैक

VACCINATION SCHEDULE

Primary Vaccination: At the age of 4 months and above; Booster Dose: 4 weeks after primary vaccination, required for complete protection. Re-vaccination: Every 6 months as per guidelines of LHDCP,

VACCINATION OF PREGNANT ANIMALS

FH VAC® can be used for vaccination of pregnant animals. However, to be on safer side, the vaccination of the animals in advanced stage of pregnancy may be avoided.

PRECAUTIONS

Immunize only healthy animals. Malnutrition, infestation with worms, administration of immunosuppressive agents such as corticosteroids or radiotherapy will interfere with the immune response to vaccine.

POST-VACCINATION REACTIONS

Generally, there are no adverse reactions noticed. Occasionally a transient palpable swelling may occur at the site of injection, which will subside within 48 to 72 hours.

STORAGE & TRANSPORTATION

Store and transport the vaccine between 2 and 8°C until use. At no stage should the vaccine be allowed to freeze. Protect from sunlight.

SHAKE WELL BEFORE USE



Manufactured by:

Brilliant Bio Pharma Private Limited
6-2-1012, TGV Mansion, 5th Floor, Khairatabad
Hyderabad - 500004 Telangana, INDIA.



Tel: +91 40 6666 7464 / 6677 2726

Email: info@bbpl.co.in

www.brilliantbiopharma.com